

# Sets

## Sets

- a set is any well-defined collection of objects
- small sets may be introduced by listing their elements
- other sets may be constructed using set comprehension, the power set operator, and the Cartesian product.

## Examples

- the four oceans of the world
- the individuals who have been appointed to the post of secretary-general of the United Nations
- the prime numbers
- the collection of programs written in  $C^{++}$  that halt if run for a sufficient time on a computer with unlimited storage

## Extension

- *Oceans* == { *Atlantic*, *Arctic*, *Indian*, *Pacific* }
- *Secretaries-General* ==  
    { *Trygve Lie*, *Dag Hammarskjöld*, *U Thant*,  
      *Kurt Waldheim*, *Javier Pérez de Cuéllar*,  
      *Boutros Boutros Ghali*, *Kofi Annan* }

## Membership

We write  $x \in s$  to indicate that  $x$  is an element of set  $s$ .

We write  $\neg(x \in s)$  as  $x \notin s$ .

## Example

- $3 \in \text{Primes}$
- $5 \in \text{Primes}$
- $8 \notin \text{Primes}$

## Membership and extension

An object  $t$  is an element of a set written in extension iff it is equal to one of the expressions within the braces:

$$\frac{t = u_1 \vee \dots \vee t = u_n}{t \in \{u_1, \dots, u_n\}} \text{ [membership and extension]}$$

## Extensionality

Two sets are equal iff every element of one is also an element of the other:

$$\frac{(\forall x : t \bullet x \in u) \wedge (\forall x : u \bullet x \in t)}{t = u} \text{ [extensionality]}$$

provided that  $x$  is free in neither  $u$  nor  $t$

## Subset

A set  $s$  is a subset of another set  $t$  iff every element of  $s$  is also an element of  $t$ :

$$\frac{\forall x : s \bullet x \in t}{s \subseteq t} \text{ [subset]}$$

provided that  $x$  is not free in  $t$

## Examples

- $\{1, 2, 3\} \subseteq \{0, 1, 2, 3, 4\}$
- $\{4, 6, 8\} \subseteq \{4, 6, 8\}$

## The empty set

$$\underline{A \times a \bullet x \notin \emptyset} \quad [\text{empty set}]$$

## The natural numbers

$$\mathbb{N} = \{0, 1, 2, 3, 4, 5, \dots\}$$

## Set comprehension

We may restrict our attention to those elements of a set  $s$  that possess a property  $p$ :

$$\begin{aligned} & \underline{\underline{t \in \{x : s \mid p\}}} \quad [\text{comprehension}] \\ & t \in s \wedge p[t/x] \end{aligned}$$

## Example

$$jim \in \{p : Person \mid p \text{ eats cheese}\}$$

## Term comprehension

We may also describe a set of objects constructed from certain elements of a given set:

$$\frac{f \in \{x : s \mid p \bullet e\}}{\exists x : s \mid p \bullet e = f} \quad [\text{term comprehension}]$$

provided that  $x$  is not free in  $f$

## Example

56, West Street  $\in$

$\{p : \text{Person} \mid p \text{ eats cheese} \bullet \text{address}(p)\}$

## Power set

$$\begin{array}{l} s \subseteq a \\ \hline s \in \mathbb{P} a \end{array} \text{ [power]}$$

## Examples

- $\mathbb{P}\{x, y\} = \{\emptyset, \{x\}, \{y\}, \{x, y\}\}$
- $\{1, 2, 3, 4\} \in \mathbb{P}\mathbb{N}$
- $\mathbb{P}\emptyset = \dots?$

## Cartesian product

If  $a$  and  $b$  are sets, then the Cartesian product  $a \times b$  consists of all tuples of the form  $(x, y)$ , where  $x$  is an element of  $a$  and  $y$  is an element of  $b$ .

$$\underbrace{x_1 \in a_1 \wedge \dots \wedge x_n \in a_n}_{(x_1, \dots, x_n) \in a_1 \times \dots \times a_n} \quad [\text{tuple membership}]$$

## Example

$$\text{Days} = \{1, 2, \dots, 31\}$$

$$\text{Months} = \{\text{January}, \text{February}, \dots, \text{December}\}$$

$$\text{Years} = \{1900, 1901, \dots, 1999\}$$

$$(22, \text{November}, 1990) \in \text{Days} \times \text{Months} \times \text{Years}$$

## Ordered tuples

Two tuples are equal if they have exactly the same elements, and these are arranged in the same order:

$$\begin{array}{l} \underline{\underline{x_1 = y_1 \wedge \dots \wedge x_n = y_n}} \\ \underline{\underline{(x_1, \dots, x_n) = (y_1, \dots, y_n)}} \end{array} \quad \text{[tuple equality]}$$

## Component selection

We may select a component of an ordered tuple using a simple dot notation: if  $m$  is an integer between 1 and  $n$ , then

$$(x_1, x_2, \dots, x_n) \cdot m = x_m$$

## Example

*date* = (22, November, 1990)

*date.1* = 22

*date.2* = November

*date.3* = 1990

## Questions

{ *x* : *Person*; *y* : *Person* | *x* is associated with *y* ∧

*x* eats cheese ∧

*y* reads Hello • *x* }

- what if we are looking for both of them?
- what if we omit the term part?

## Characteristic tuple

The expression

$$\{ a : \mathbb{N}; s : \mathbb{P}\mathbb{N}; t : \mathbb{P}\mathbb{N} \mid a \in s \wedge a \notin t \}$$

has the same value as

$$\{ a : \mathbb{N}; s : \mathbb{P}\mathbb{N}; t : \mathbb{P}\mathbb{N} \mid a \in s \wedge a \notin t \bullet (a, s, t) \}$$

## Question

Can you exhibit an element of each of the following sets?

- $\{ a : \mathbb{N}; b : \mathbb{N} \mid a > b \}$
- $\{ s : \mathbb{P}\mathbb{N}; a : \mathbb{N}; t : \mathbb{P}\mathbb{N} \mid a \in s \wedge a \in t \}$

**Union**

$$\frac{x \in (a \cup b)}{x \in a \vee x \in b} \text{ [union]}$$

**Example**

$$\begin{array}{l} \text{Guests} = \text{Jim's friends} \\ \cup \\ \text{Eleanor's friends} \end{array}$$

**Intersection**

$$\frac{x \in (a \cap b)}{x \in a \wedge x \in b} \text{ [intersection]}$$

**Example**

$$\text{Steve} \in \text{Choir} \cap \text{1st VIII} \cap \text{1st XV}$$

## Difference

$$\frac{x \in (a \setminus b)}{x \in a \wedge x \notin b} \quad [\text{diff}]$$

## Example

$$Jim \in Voters \setminus Conservatives$$

## Distributed forms

$$\frac{x \in \bigcup s}{\exists a : s \bullet x \in a} \quad [\text{distributed union}]$$

$$\frac{x \in \bigcap s}{\forall a : s \bullet x \in a} \quad [\text{distributed intersection}]$$

## Example

$$\text{Europeans} = \bigcup \{ \text{Belgians, British, Danes, Dutch,} \\ \text{French, Germans, Greeks, Irish,} \\ \text{Italians, Luxemburgers, Portuguese,} \\ \text{Spanish, Swedes, \dots} \}$$

$$\{\text{Steve}\} = \bigcap \{ \text{Choir, 1st VIII, 1st XV} \}$$

## Types

A **type** is a maximal set, at least within the confines of the current specification.

This ensures that each object **x** is associated with exactly one type: the largest set **s** present for which  $x \in s$ .

## Integers

There is a single built in type,  $\mathbb{Z}$ , the type of all integers

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Other types may be constructed from  $\mathbb{Z}$ , or from user-defined basic types.

## Examples

- $\mathbb{Z} \times \mathbb{Z}$
- $\mathbb{P} \mathbb{Z}$
- *People*
- ...

## Advantages

- adds structure and readability
- correctness via type-checking
- data types suggest implementation

**Question**

What is the type of the empty set?

**Summary**

- collections of objects
- membership
- extensionality
- comprehension
- $\mathbb{P}$  and  $\times$
- types